

Turf Fertilizer with

INSECTICIDE

GROUP

LEPRYN® Ш .067% ACI

Counties of New York State. recreational turfgrass and sod farms and Suffolk, Kings, Queens as well as landscape ornamentals, interior plantscapes infesting landscape in Nassau, and other listed pests Sale Into, Distribution and/or Use grubs white (including golf courses) control of Not for Sale, systemic For

Analysis Guaranteed

Total Nitrogen (N)17.0%	17.0%
12,7% Urea Nitrogen†	
1.9% Ammoniacal Nitrogen	
2.3% Water Insoluble Nitrogen (W.I.N.)*	
0.1% Other Water Soluble Nitrogen*	
Soluble Potash (K ₂ 0)4.0%	%0:
Calcium (Ca)2.4%	2.4%
Sulfur (S)2.1%	2.1%
2.1% Combined Sulfur	
Iron (Fe)0.48%	0.48%
0.05% Water Soluble Iron	
Chlorine (CI) not more than2.9%	%6

diamide chemical class

anthranilic

Chlorantraniliprole belongs to the

and Coverage

Rates

Application

Suggested

100.000%

...0.067%

ide

-1-(3- chloro-2-pyridinyl)-1H-pyrazole-5-carboxar

Active Ingredient

Chlorantraniliprole*

Other Ingredients

Total.....

88 lbs/K

Ø

44lbs/K 58#N/K

ന

sq.ft.

,534

4,

COVERAG

16#N/K

က

E

LESCO Letter/Number Scotts R-8

X

Rate Rate

G/16 M M

J 2

Prizelawn HVN - Earthway

 $|\infty|$

4 2 2

Spyker Lely

Derived From

Muriate of potash, Polymer-coated Biosolids, Ammonium : urea, Urea

6.1% Slowly

by Elmsford, NY 10523 Phone # 914-347-5656 Net Weight 50 Pounds Guaranteed & Irrigation 8 Williams St. and Central Turf Distributed

Product Code:

Premium Turf 0124 Rev 220

Grade: Lot#

14921 SGN

5. NO. 100-1492-93243 . NO. 93243-CT-1 (CT) . NO. 90561-NC-001 (N) . NO. 93243-TX-1 (TX)





Hold eye open and rinse slowly and gently with water for 15-20 minutes Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment, HOT LINE NUMBER

Precautionary Statements

Keep Out of Reach of Children

CAUTION

CAUTION

Hazards to Humans and Domestic Animals

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

For 24-Hour Medical Emergency Assistance (Human or Animal), or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident) Call 1-800-331-3148

- PERSONAL PROTÉCTIVE EQUIPMENT (PPE) Applicators and other handlers must wear
- Long-sleeved shirt and long pants

IF IN EYES:

After the product has been loaded into properly calibrated application equipment, shirt, pants, socks, and shoes are sufficient Personal Protective Equipment. Follow manufacturer's instructions for cleaning/ maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry

User Safety Recommendations

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 Remove clothing immediately if pesticide gets inside.
- . Then wash thoroughly and put on clean clothing.

Environmental Hazards

Hudson Valley Turf Fertilizer with 0.067% Acelepryn Insecticide Herbicide is toxic to aquatic invertebrates, oysters and shrimp. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to use sites,

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read and understand the entire label before using this product.

Hudson Valley Turf Fertilizer with 0.067% Acelepryn Insecticide must be used only in accordance with specifications on this label or in separate Central Turf & Irrigation supplemental labeling that may be made temporarily available through local distributors, as a result of new EPA approvals. Central Turf & Irrigation will not be responsible for losses or damages resulting from the use of this product in any manner not specifically stated on this label or other labels or bulletins published by Central Turf & Irrigation. User assumes all risks associated with such non-specified use.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment, restricted-entry interval, and notification to workers (as applicable).

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours. For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear: Long-sleeved shirt and long pants

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides, 40 CFR part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Professional applications to golf courses, residential, industrial and commercial lawns and sports fields are not within the scope of the Worker Protection Standard

Do not allow other workers to enter the treated area while application is in progress. Do not apply this product in commercial nurseries and greenhouse

Product Information

Hudson Valley Turf Fertilizer with 0.067% Acelepryn Insecticide may be applied for the control of white grubs and other listed pests infesting landscape and recreational turfgrass (including golf courses) as well as

Hudson Valley Turf Fertilizer with 0.067% Acelepryn Insecticide may be used on plants or turfgrasses that are being grown for aesthetic or recreational purposes or climatic modification in or around home lawns, commercial lawns, industrial facilities, residential dwellings, business and office complexes, shopping complexes, multi-family residential complexes, institutional buildings, airports, cemeteries, interior plantscapes, ornamental gardens, parks, wildlife plantings, playgrounds, schools, day-care facilities, golf courses (tee box areas, roughs, fairways, greens, collars etc.), athletic fields, sod farms and other landscaped areas. Do not apply Hudson Valley Turf Fertilizer with 0.067% Acelepryn Insecticide when the soil is saturated with water because adequate distribution of the active ingredient vertically in the soil profile cannot be

Hudson Valley Turf Fertilizer with 0.067% Acelepryn Insecticide is in GROUP 28 of the EPA's Insecticide and Acaricide Groups Based on Target Site of Action (EPA PR Notice 2001-5) and may be used in rotational resistance management programs

Consult your local Central Turf & Irrigation representative, Cooperative Extension Service specialist or pest control advisor for regionally specific information regarding application timing.

Apply Hudson Valley Turf Fertilizer with 0.067% Aceleptyn Insecticide with drop-type, rotary-type or hand-held application equipment. The application equipment must be properly calibrated according to the manufacturer's directions to ensure uniform coverage over the treated area. During application, check equipment periodically to confirm that it is working properly and that the granules are being uniformly distributed.

APPLICATION TIMING

White Grubs
Apply Hudson Valley Turf Fertilizer with 0.067% Acelepryn Insecticide from early April to early September for preventative and early curative control all of the major white grub species infesting turfgrass. The need for an application may be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. The higher rate listed for white grub control in Table 1 may be required for early season (April and May) applications where a long residual is required or in late August or early September when less sensitive mid-instar grubs are present at the time of application. Irrigate turf immediately after

Annual Bluegrass Weevil
Apply Hudson Valley Turf Fertilizer with 0.067% Aceleptyn Insecticide when overwintered adult annual bluegrass weevils are observed in late April or early May to prevent damage from first generation larvae in late-May and June. An application of Hudson Valley Turf Fertilizer with 0.067% Aceleptyn e at this time will also provide white grub control. The higher rates listed in Table 1 may be required for applications made after mid-May.

Annly Hildson Valley Turf Fertilizer with 0.067% Acelebryn Insecticide when overwintered adult billbugs are first observed. This will usually occur in late April or early May in regions with cool-season turfgrasses. An application of Hudson Valley Turf Fertilizer with 0.067% Acelepryn Insecticide at this time will also provide white grub control. The higher rates listed in Table 1 may be required for applications made after early May or when billbug species other than bluegrass billbug are present.

European Crane Fly
Apply Hudson Valley Turf Fertilizer with 0.067% Acelepryn Insecticide between September and November to control European crane fly larvae in turfgrass. The higher rate listed in Table 1 may be required to achieve control when applications are made in November.

Spittlebugs
Hudson Valley Turf Fertilizer with 0.067% Acelepryn Insecticide will provide control of two-lined spittlebug when applications are made in spring or

Turf Caterpillars

Hudson Valley Turf Fertilizer with 0.067% Acelepryn Insecticide will provide curative caterpillar control in turfgrass. If the area being treated is maintained at a mowing height of greater than one inch, then the higher rate listed in Table 1 may be required during periods of high pest pressure

For suppression of chinch bugs, apply Hudson Valley Turf Fertilizer with 0.067% Acelepryn Insecticide before eggs hatch.

APPLICATION RATES FOR LAWNS, GOLF COURSES AND OTHER RECREATIONAL TURFGRASS AREAS

Table 1 provides the application rates that will control the listed turfgrass pests. For maximum residual effectiveness or for optimal control of large pest infestations, apply Hudson Valley Turf Fertilizer with 0.067% Acelepryn Insecticide at up to 500 pounds per acre (0.33 lb Al/A) to control any of the pests listed in Table 1. Restriction: Do not apply more than 750 pounds (equivalent to 0.5 lb of active ingredient) of product per acre per year in broadcast applications to turfgrass.

TABLE 1: TURFGRASS APPLICATION RATES

Target Pest	Product per Acre	Product per 1,000 sq ft	lb Al/A
White Grubs (including Aphodius spp., Asiatic garden beetle, black turfgrass ataenius, European chafer, green June beetle, Japanese beetle, May/June beetles (Phyllophaga spp.), northern masked chafer, oriental beetle and southern masked chafer) Billbugs European Crane Fly	150–300 lb*	3.44 – 6.88 lb	0.1–0.2
Turf caterpillars (including armyworms, cutworms and sod webworms)			
Annual bluegrass weevil	225–300	5 17–6 88 lb	0.15-0.2
Spittlebugs	lb*	5.17 – 5.56 lb	0.13-0.2
Chinch bugs (suppression only)	150–300 lb*	3.44–6.88 lb	0.1–0.2

^{*}For maximum residual effectiveness or optimal control of large pest infestations, apply Hudson Valley Turf Fertilizer with 0.067% Acelepryn Insecticide at up to 500 pounds per acre (0.33 lb Al/A).

APPLICATION RATES FOR ORNAMENTAL PLANTS

Table 2 provides the application rates that will control the pests of landscape ornamentals and interior plantscapes. Before application, remove black plastic or other weed barrier materials that are impermeable to water. For maximum residual effectiveness or for optimal control of large pest infestations, Hudson Valley Turf Fertilizer with 0.067% Aceleptyn Insecticide may be applied at up to 500 pounds per acre (0.33 lb Al/A) to control any of

Restriction: Do not apply more than 750 pounds (equivalent to 0.5 lb of active ingredient) of product per acre per year in broadcast applications to

TABLE 2: ORNAMENTAL TREE AND SHRUB (INCLUDING EVERGREENS), FLOWER, FOLIAGE PLANT, AND GROUNDCOVER APPLICA-

Target Pest	Product per Acre	Product per 1,000 sq ft	lb Al/A
White Grubs (Asiatic garden beetle, European chafer, green June beetle, Japanese beetle, May/June beetles (Phyllophaga spp.), northern masked chafer, oriental beetle and southern masked chafer)	150–300 lb*	3.44 – 6.88 lb	0.1–0.2

*For maximum residual effectiveness or optimal control of large pest infestations, apply Hudson Valley Turf Fertilizer with 0.067% Acelepryn Insecticide at up to 500 pounds per acre (0.33 lb Al/A).

We recommend that you follow the Florida Green Industries BMPs at: http://www.flaes.org/pdf/BMP_Book_final.pdf We recommend that you follow the Florida Golf Course BMPs at: http://www.flaes.org/pdf/glfbmp07.pdf

If this product is being sold in a state requiring the publication of information regarding its metal content, information regarding the contents and levels of metals in this product is available at the following website: http://aapfco.org/metals.htm, or by calling 1-877-220-1014.

CLEANING APPLICATION EQUIPMENT

Thoroughly clean all application equipment after use to reduce the risk of forming hardened deposits that might become difficult to remove. Thoroughly rinse application equipment with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean equipment near wells, or other water sources. Dispose of waste rinse water in accordance with local regulations.

INTEGRATED PEST MANAGEMENT (IPM) PROGRAMS

Hudson Valley Turf Fertilizer with 0.067% Acelepryn Insecticide is recommended for IPM programs on turf and landscape ornamentals because it does not directly impact natural arthropod predator and parasitoid populations including ladybird beetles, lacewings, minute pirate bugs and predatory mites. The feeding behavior of beneficial arthropods will aid in extending natural control of other insect and mite pests and will reduce the possibility of secondary pest outbreaks. Hudson Valley Turf Fertilizer with 0.067% Acelepryn Insecticide will reduce the target pest species that may serve as a food source for beneficial arthropods

RESISTANCE MANAGEMENT

Some insects are known to develop resistance to products used repeatedly for control. When this occurs, the labeled dosages fail to suppress the pest population below threshold levels. Because the development of resistance cannot be predicted, the use of this

product should conform to resistance management strategies established for the use area.

These strategies may include incorporation of cultural and biological control practices, alternation of active classes of insecticides on succeeding generations and targeting the most susceptible life stage. Consult your local Central Turf & Irrigation representative Cooperative Extension Service specialist or pest control advisor for the latest information on resistance management in your area.

- · Not for use on plants (except turfgrass on sod farms) being grown for sale or other commercial use, or for commercial seed
- Do not formulate this product into other end-use products without written permission from Central Turf & Irrigation .
- Do not apply this product incomercial nurseries and greenhouses.
 Do not apply this product in commercial nurseries and greenhouses.
 Do not apply more than 750 pounds (equivalent to 0.5 lb of active ingredient) of product per acre per year in broadcast applications to turfgrass or landscape ornamental plants.

 • Do not apply more than 300 pounds (equivalent to 0.2 lb of active ingredient) of product per acre in a single application on sod
- Wait a minimum of 7 days to retreat.
- · Do not apply by air.

The following restrictions are required to permit use of Hudson Valley Turf Fertilizer with 0.067% Acelepryn Insecticide in the State of New York: Golf courses

Greens, tee boxes: Do not apply this product within 25 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch). Fairways: Do not apply this product within 50 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

All other use sites: Do not apply this product within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal

Store product in original container only in a location inaccessible to children and pets. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Not for use or storage in or around the home.

Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Central Turf & Irrigation or Seller. To the extent permitted by applicable law, Buyer and User agree to hold Central Turf & Irrigation and Seller harmless for any claims relating to such factors.

Central Turf & Irrigation warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or Central Turf & Irrigation , and, (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW. Central Turf & Irrigation MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall Central Turf & Irrigation be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE LIABILITY OF CENTRAL TURF & IRRIGATION AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CENTRAL TURF & IRRIGATION OR SELLER. THE REPLACEMENT OF THE PRODUCT.

Central Turf & Irrigation and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Central Turf & Irrigation.

Acelepryn is a Trademark of a Syngenta Group Company

*Do not use on food producing trees, vines or plants.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a dry place. Store in locked area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal

CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. Empty container through use according to label instructions. Offer for recycling, if available. If not available, then dispose of empty bag in a sanitary landfill or by other procedures allowed by state and local authorities.

^{**}Not for use on container grown plants.